

Quark distributions in QCD sum rules: unexpected features and paradoxes

A.G.Oganesian

Institute of Theoretical and Experimental Physics
B.Chermushkinskaya 25, 117218, Moscow, Russia

Abstract

Some very unusual features of the hadron structure functions, obtained in the generalized QCD sum rules, like the surprisingly strong difference between longitudinally and transversally polarized ρ mesons structure functions and the strong suppression of the gluon sea in longitudinally polarized ρ mesons are discussed. Also the problem of exact zero contribution of gluon condensates to pion and longitudinally polarized ρ meson quark distributions is discussed.

It is well known, that in QCD DGLAP equations predicts the evolution of these distributions with Q^2 but not the initial values at some relatively low Q^2 from which this evolution starts. To obtain these initial values (inputs) usually any form of quark (valence and sea) and gluon distributions with some free parameters are assumed. Then by using DGLAP equations, quark and gluon distributions are calculated at all Q^2 and x and compared with the whole set of the experimental data, and in this way the parameters are fixed. Evidently, such an approach is not completely satisfactory from theoretical point of view, because, we can loose some unexpected physical features of distributions, so it would be desirable to determine the initial distribution directly from QCD. This talk will be devoted to the problem of the determination of quark and gluon distributions in hadrons in model independent way in QCD, and we will see, that some unexpected features of quark distributions really appear.

A method to determinate valence quark distribution in QCD sum rules at intermediate x was suggested in [1] and generalized in [2]-[3]. Due to this generalization it became possible to calculate valence quark distributions in pion [2] and transversally and longitudinally polarized ρ -meson [3], and also significantly improve the result for valence quark distributions in proton [4].

Let me at first briefly remind the method. We start from consideration of the 4-point correlator corresponding to the forward scattering of hadron and electromagnetic currents.

$$\Pi = -i \int d^4x d^4y d^4z e^{ip_1x + iqy - ip_2z} \langle 0 | T \{ j^h(x), j^{el}(y), j^{el}(0), j^h(z) \} | 0 \rangle \quad (1)$$

Here p_1 and p_2 are the initial and final momenta carried by hadronic the current j^h , q and $q' = q + p_1 - p_2$ are the initial and final momenta carried by virtual photons (Lorentz indices are omitted) and $t = (p_1 - p_2)^2 = 0$.

The idea of the approach (in the improved version) is to consider the imaginary part (in s -channel) of this four-point correlator $\Pi(p_1, p_2, q, q')$. It is supposed that virtualities of the photon q^2, q'^2 and hadron currents p_1^2, p_2^2 are large and negative $|q^2| = |q'^2| \gg |p_1^2|, |p_2^2| \gg R_c^{-2}$, where R_c is the confinement radius. It was shown in [1] that in this case the imaginary part in s -channel [$s = (p_1 + q)^2$] of $\Pi(p_1, p_2; q_1, q')$ is dominated by a small distance contribution at not small x , so OPE is applicable.

To find structure function one should compare dispersion representation of the forward scattering amplitude in terms of physical states with those in OPE and use Borel transformation. Though we should consider the case of forward scattering, i.e. $p_1 = p_2$, but, and it is significant point of the method, we should keep p_1^2 not equal to p_2^2 in all intermediate stages and only in final result take the forward scattering limit. Only in such way it is possible to effectively (exponentially) suppress all terms in sum rules except those which are proportional to structure function and achieve good accuracy. Finally equating the physical and QCD representations (for detail see [2]) after double borelization (on p_1^2 and p_2^2) we found

$$Im \Pi_{QCD}^0 + \text{Power correction} = 2\pi F_2(x, Q^2) g_h^2 e^{-m_h^2(\frac{1}{M_1^2} + \frac{1}{M_2^2})} \quad (2)$$

Here $Im \Pi_{QCD}^0$ is the bare loop contribution, (where continuum contribution is eliminated) and g_h is defined as $\langle 0 | j_h | h \rangle = g_h$. In what follows, we will suppose $M_1^2 = M_2^2 \equiv 2M^2$.

Let us start with the case of the pion. To find the pion structure function one should choose the imaginary part of 4-point correlator with two axial (hadron) and two electromagnetic currents and consider the invariant amplitude at tensor structure, $P_\mu P_\nu P_\lambda P_\sigma / \nu$, where $P = (p_1 + p_2)/2$, μ, ν are vector current indexes and λ, σ -hadron current indexes. I shall briefly note the main points of calculations, for detail see [2]. In QCD part of sum rules we take into account the following terms:

1. Unit operator contribution (bare loop, Fig.1) and leading order (LO) perturbative corrections proportional to $\ln(Q^2/\mu^2)$, where μ^2 is the normalization point. In what follows, the normalization point will be chosen to be equal to the Borel parameter $\mu^2 = M_1^2 = M_2^2 = 2M^2$.

2. Power corrections – higher order terms of OPE. Among them, first of all we should account the dimension-4 correction, proportional to the gluon condensate $\langle 0 | \frac{\alpha_s}{\pi} G_{\mu\nu}^n G_{\mu\nu}^n | 0 \rangle$. Corresponding diagrams are shown on Fig.2. But, surprisingly, it was found that these diagrams contribution to the sum rule is exactly cancelled after Borel transformation on p_1^2 and p_2^2 . I want to note, that diagrams in total may not cancel, but only that part of them, which have imaginary part both p_1^2 and p_2^2 (i.e. just those which contribute to the pion structure function). So we see there is no gluon condensate contribution to the sum rule for pion structure function. The fact of exact cancellation of contributions of 10 diagrams on fig.2 is surprising, but of course one can treat this as a just accident. But just the same take place for operators of the next dimension!

There are a large number of loop diagrams, corresponding to $d = 6$ corrections. First of all, there are diagrams which correspond to interaction only with gluon vacuum field, i.e. only with external soft gluon lines (see Fig.3). They are, obviously, proportional to $\langle g^3 f^{abc} G_{\mu\nu}^a G_{\alpha\beta}^b G_{\rho\sigma}^c \rangle$. But one should also take into account diagrams of Fig.4 which are

proportional to $\langle 0 | D_\rho G_{\mu\nu}^a D_\tau G_{\alpha\beta}^a | 0 \rangle$ and $\langle 0 | G_{\mu\nu}^a D_\rho D_\tau G_{\alpha\beta}^a | 0 \rangle$. Using the equation of motion it can be expressed in the terms of

$$[\langle 0 | g\bar{\psi}\psi | 0 \rangle]^2 \quad \text{and} \quad \langle 0 | g^3 G_{\mu\nu}^a G_{\nu\rho}^b G_{\rho\mu}^c f^{abc} | 0 \rangle$$

As it was shown in [2], if one take the sum of all diagrams, then the terms, proportional to gluon condensate $\langle 0 | g^3 G_{\mu\nu}^a G_{\nu\rho}^b G_{\rho\mu}^c f^{abc} | 0 \rangle$ exactly cancel each other just in the same way, as for $d = 4$ operator we have discussed above. I want to note, that there is about a hundred different diagrams, so it is hard to believe that such a cancellation is again an accident.

So we can fixed first very unexpected feature of pion dynamics - gluon vacuum contribution to pion valence quark distribution is exact zero for dimensions 4,6. I want to add, that, as we will see a little later, just the same effect appear for longitudinally polarized ρ meson, so this is not the specific feature of pion. It is very like that here we meet a consequence of some symmetry, though today we cannot understand it.

Of course except the diagrams on fig. 2a,b there are another diagrams of dimension 6, directly proportional to the four-quark operators $a^2 = \alpha_s(2\pi)^4(\langle 0|\bar{\psi}\psi|0\rangle)^2$. I will not discuss them in this talk, one can see papers [2], [3]. Finally quark distribution function (hereafter we will note it as QDF) has the following form:

$$\begin{aligned} xu_\pi(x) = & \frac{3}{2\pi^2} \frac{M^2}{f_\pi^2} x^2(1-x) \left[\left(1 + \left(\frac{a_s(\mu^2) \cdot \ln(Q_0^2/\mu^2)}{3\pi} \right) \right. \right. \\ & \times N(x)(1 - e^{-s_0/M^2}) - \left. \left. \frac{4\pi\alpha_s(\mu^2) \cdot 4\pi a^2}{(2\pi)^4 \cdot 3^7 \cdot 2^6 \cdot M^6} \cdot \frac{\omega(x)}{x^3(1-x)^3} \right] \end{aligned} \quad (3)$$

where $N(x) = (1-x)(1+4x\ln(1-x)) - 2x(1-2x)\ln x$ and $\omega(x)$ is the fourth degree polynomial in x , This function $u_\pi(x)$ may be used as an initial condition at $Q^2 = Q_0^2$ for solution of QCD evolution equations. The result is shown on fig.5 (thick line).

In the numerical calculations we choose $Q_0^2 = 2 \text{ GeV}^2$ (for pion) and $a^2 = 0.23 \text{ GeV}^6$, found recently from τ decays analysis -see [5], [6], [7]. The continuum threshold was varied in the interval $0.8 < s_0 < 1.2 \text{ GeV}^2$ and it was found, that the results depend only slightly on it's variation. The analysis of the sum rule shows, that it is fulfilled in the region $0.2 < x < 0.7$, where both power correction and continuum contribution are less than 30% and the stability in the Borel mass parameter M^2 dependence in the region $0.4 \text{ GeV}^2 < M^2 < 0.7 \text{ GeV}^2$ is good. The accuracy of this result was estimated in [2] to be 20 – 30%. Comparison with various experimental result (see [2]) lead to conclusion, that agreement with experiment is good. I want to note, that this result for QDF is based only on the QCD sum rules.

It is possible also to estimate the second moments of QDF. Assume, that $u_\pi(x) \sim 1/\sqrt{x}$ at small $x \lesssim 0.15$ according to the Regge behavior and $u_\pi(x) \sim (1-x)^2$ at large $x \gtrsim 0.7$ according to quark counting rules. Then, matching these functions with our result, one may find the numerical values of the first and the second moments of the u -quark distribution: $\mathcal{M}_1 = \int_0^1 u_\pi(x) dx \approx 0.84$, $\mathcal{M}_2 = \int_0^1 xu_\pi(x) dx \approx 0.21$ where the values various slightly at reasonable changes of QDF behavior at large or small x . The moment $\mathcal{M}_1$ has the meaning of the number of u quarks in π^+ and it should be $\mathcal{M}_1 = 1$. The

deviation of M_1 from 1 characterizes the accuracy of our calculation. The moment $\mathcal{M}_2$ has the meaning of the pion momentum fraction carried by the valence u quark. Therefore, the valence u and $\bar{d}$ quarks carry about 40% of the total momentum what is close to experimental results.

In the same way one can calculate valence u -quark distribution in the ρ^+ meson. The choice of hadronic current is evident. $j_\mu^\rho = \bar{u}\gamma_\mu d$ and the matrix element is $\langle \rho^+ | j_\mu^\rho | 0 \rangle = \frac{m_\rho^2}{g_\rho} e_\mu$ where m_ρ is the ρ -meson mass, g_ρ is the $\rho - \gamma$ coupling constant, $g_\rho^2/4\pi = 1.27$ and e_μ is the ρ meson polarization vector.

In the non-forward amplitude the tensor structure for determination of u -quark distribution in the longitudinal ρ meson is to $P_\mu P_\nu P_\sigma P_\lambda$, while for transverse ρ it is $-P_\mu P_\nu \delta_{\lambda\sigma}$ (see [3]).

In the case of longitudinal ρ meson the tensor structure, that is separated is the same as in the case of the pion. It was shown in [3], that u -quark distribution in the longitudinal ρ meson can be found from Eq.(3) by substituting $m_\pi \rightarrow m_\rho$, $f_\pi \rightarrow m_\rho/g_\rho$. Sum rules for $u_\rho^L(x)$ are satisfied in the wide x region: $0.1 < x < 0.85$ with high accuracy (about 10%). Fig.5 (curve with squares) presents $xu_\rho^L(x)$ as a function of x . The values $M^2 = 1 \text{ GeV}^2$ and $s_0 = 1.5 \text{ GeV}^2$, $Q_0^2 = 4 \text{ GeV}^2$ were chosen. It is clear, that in this case again gluon condensate contribution is cancelled. If we construct the moment of QDF for ρ_L in the same way as for pion then we found $M_1 = 1.06$ and $M_2 = 0.39$ (for one valence quark). One should note, that accuracy of this prediction of moment of ρ_L is very high, because in this case sum rules for QDF are covered almost all x region, as I have noted before, so extrapolation procedure contribution numerically is negligible.

So we can fix the second very unusual effect: the valence quarks carry about 80% of the total momentum of longitudinally polarized ρ meson, so gluon sea should be strongly suppressed (less than 20%).

Let us now consider the case of transverse ρ -meson, i.e., the term proportional to the structure $P_\mu P_\nu \delta_{\lambda\sigma}$. The procedure of calculations are the same except one significant points.

1. In contrast to the pion case, the $\langle G_{\mu\nu}^a G_{\mu\nu}^a \rangle$ correction for transversally polarized $\rho(\rho_T)$, .

2. In contrast to π and ρ_L -meson cases, the terms proportional to $\langle G_{\mu\nu}^a G_{\mu\nu}^a \rangle$, as long as $\langle 0 | g^3 f^{abc} G_{\mu\nu}^a G_{\nu\rho}^b G_{\rho\mu}^c | 0 \rangle$ are not cancelled for ρ_T . But $\langle 0 | g^3 G_{\mu\nu}^a G_{\nu\rho}^b G_{\rho\mu}^c f^{abc} | 0 \rangle$ is not well known; so we need to use estimations based there are only some estimates based on the instanton model. Result for QDF in transversally polarized ρ meson is shown on Fig.5 for the region $0.2 < x < 0.65$, (line with asters). One can find complete analytical form and it detail analysis in our paper [3]. The choose of parameters are the same as in previous cases. The main sources of inaccuracy are $d = 6$ gluon condensate (gives about 20%) and $d = 4$ gluon condensate (due the inaccuracy of it own value about factor 1.5, which lead to uncertainty in QDF about 20%). Accuracy is worse, about 30 – 50% (better in the middle of region and worse at the end). From fig.5 one can see, that difference between QDF in the longitudinally (ρ_L) and transversally (ρ_T) polarized ρ mesons is very large, many times larger that uncertainties.

So, and this is third surprise, we can conclude that QDF significantly depend of polarization. Such great difference between QDF of ρ_L and ρ_T is very difficult to explain

in usual quark model. I want to add, that to check this effect, we calculate the second moment of QDF of ρ_L and ρ_T in quite another way, considering of 2-point correlator in external tensor field. This method was offered in [8] for pion case many years ago (for the method itself of the QCD sum rules in the external fields see [9]). In the [10] we use this method for polarized ρ meson case. The results totally confirm the conclusion that QDF of ρ_L and ρ_T are quite differ : the second moment M_2 for one ρ_L is about 84% (compare with 78% discussed above) while M_2 for ρ_T is much smaller and close to usual hadron case (about 40% - 50%). (More detail this results and comparison of this two methods one can found in [10].) That why I think, that QDF indeed strongly depend on polarization, at least in ρ -meson case, and this fact can't be explained in the present quark models of the hadrons.

Summarizing, in this talk I tried to pay your attention to very strange results, which appear from pure QCD non-model calculation of quark distribution function and yet have no satisfactory physical explanation. These are:

a) First of all, the total cancellation of diagrams with gluon condensate (for dimensions $d = 4, 6$) seems to note to some unknown symmetry. Of course it is interesting to check this property also for dimension $d = 8$, but it is technically very hard and isn't complete yet.

b) Second, such a strong difference between QDF for longitudinally polarized (ρ_L) and transversally polarized (ρ_T) , as is shown on fig.5, can not be explained in the usual hadron quark models.

Taking in mind also the third very strange fact that, as we discuss before, for ρ_L gluon sea is strongly suppressed (less than 15%), we can conclude quark dynamic in (ρ_L) and (ρ_T) should be quite different, in contradiction to usual quark models. From my point of view it is extremely interesting to understand this strange features, at least in some model way.

Author is thankful to B.L.Ioffe for useful discussions.

This work was supported in part by CRDF grant RUP2-2621-MO-04 and RFFI grant 06-02-16905.

References

- [1] V.M.Belyaev and B.L.Ioffe, Nucl. Phys. B 310, (1988) 548.
- [2] B.L.Ioffe and A.G.Oganesian, Eur. Phys. J. C 13 (2000) 485.
- [3] B.L.Ioffe and A.G.Oganesian, Phys. Rev. D 63 (2001) 096006.
- [4] B.L.Ioffe and A.Oganesian, Nucl. Phys. A 714 (2003) 145.
- [5] B.L.Ioffe and K.N.Zyablyuk, Nucl. Phys. A 687 (2001) 437.
- [6] B.V.Geshkenbein, B.L.Ioffe and K.N.Zyablyuk, Phys. Rev. D 64 (2001) 093009.
- [7] B.L.Ioffe Prog.Part.Nucl.Phys. 56 (2006) 232.
- [8] V.Belyaev and B.Blok, Phys.Lett. B 167 (1986) 99, Yad.Fiz. 43 (1986) 706.

- [9] B.L.Ioffe and A.V.Smilga, Nucl.Phys. B 232 (1984), 109.
- [10] A.Oganesian and A.Samsonov, JHEP 0109: 002, (2001).




Figure 1: diagrams, corresponding to bare loop contribution




Figure 2: diagrams, corresponding to $d = 4$ gluon condensate




Figure 3: diagrams, corresponding to $d = 6$ condensate




Figure 4: diagrams, corresponding to $d = 6$ condensate




Figure 5: Quark distribution function for pion, transversally and longitudinally polarized ρ - solid line, line with asters, and line with squares correspondingly